

The Theory of Light Scattering by Semiconductor Quantum Dots. Semiclassical Method Using Retarded Potentials

I. G. Lang, L. I. Korovin

A. F. Ioffe Physical-Technical Institute, Russian Academy of Sciences, 194021 St. Petersburg, Russia

S. T. Pavlov^{†‡}

[†]*Facultad de Física de la UAZ, Apartado Postal C-580,
98060 Zacatecas, Zac., Mexico;*

E-mail: pavlov@ahobon.reduaz.mx

[‡]*P.N. Lebedev Physical Institute of Russian Academy of Sciences. 119991,
Moscow, Russia;
pavlov@sci.lebedev.ru*

The theory of elastic light scattering by semiconductor quantum dots is suggested. The semiclassical method, applying retarded potentials to avoid the problem of boundary conditions for electric and magnetic field, is used. The exact results for the Poynting vector on large distances from a quantum dot, formulas of differential cross sections of light scattering for the monochromatic and pulse irradiation are obtained.

PACS numbers: 78.47. + p, 78.66.-w

I. INTRODUCTION

The excitonic excitations in size-quantized semiconductor objects (quantum wells, quantum wires, quantum dots) may be investigated by measuring of elastic light scattering by object mentioned above. Light scattering increases resonantly when the stimulating light frequency ω_ℓ equals to the exciton frequency ω_0 . The width of the resonant peak is determined by the exciton damping Γ .

It was shown¹ that $\Gamma = \gamma_r + \gamma$, where $\gamma_r(\gamma)$ is the radiative (nonradiative) exciton damping. The same concept was extended on light absorption by quantum wells² (see also³). Light reflection by quantum wells, quantum wires, quantum dots was considered in⁴.

There are two ways to investigate theoretically light scattering by semiconductor quantum objects: quantum and semiclassical. The quantum way applies the quantum perturbation theory. The electric field is being quantized, processes of photon annihilation with an exciton creation and vice versa are considered. The quantum way was used for calculations of a light scattering section by any quantum dot. The quantum perturbation theory allows to obtain results in the lowest order on the electron-photon interaction, what excludes taking into account of the radiative damping γ_r . The nonradiative damping γ does not appear in the quantum theory too, and the resonant denominator has a form $(\omega_\ell - \omega_0)^2 + \delta^2$, $\delta \rightarrow 0$, instead of the precise expression $(\omega_\ell - \omega_0)^2 + \Gamma^2/4$. The advantage of the quantum theory is its simplicity.

With the help of the semiclassical method, classical electric and magnetic fields are calculated, whereas the description of the electron system remains quantum one (thus, there are essential the nondiagonal matrix elements of the quasimomentum operator $\mathbf{p}_{cv}$, corresponding to the electron-hole pair creation). The

semiclassical method starts since the calculation of the current and charge densities averaged on the crystal ground state¹⁶ and induced by the electric field inside of the semiconductor object^{6,7}. When the stimulating electric field $\mathbf{E}_0(\mathbf{r}, t)$ is used in expressions for these densities, we are limited again by the lowest order on the electron-light interaction. All the orders on the electron-light interaction are taken into account, when the genuine electric field is substituted into these densities.

Further, two versions of the semiclassical method may be applied. The first version is a solution of the Maxwell equations inside and outside of an object and application of the boundary conditions for electric and magnetic fields on borders of the object. This version is convenient when the refraction coefficients inside ν and outside ν_0 of the object are different. In such a way the problem of light reflection and absorption by wide quantum wells (the width of which may be comparable with the light wave length)^{8,10?}. The electric fields arising in the resonant light scattering by an exciton in a spherical quantum dot (consisting of a cubical crystal of T_d class (for example, GaAs) and limited by the infinitely rectangular potential barrier under condition $\nu_0 \neq \nu$) are calculated¹¹ by similar way. However, in the case of spherical quantum dots, application of the boundary conditions results into extremely bulky calculations, and in cases of quantum dots of other forms become even more complicated.

Therefore, we suggest a second version of the semiclassical method - the method of the retarded potential. The essence of the method is as follows. We substitute the values of the averaged induced densities of currents and charges into expressions for the vector and scalar potentials (see (7) and (8) below). Having used potentials, containing the genuine electric field inside and electric and magnetic field outside of the object, we obtain an integral equation for the electric field

inside of the object, which may be solved in some cases. Further, we determine the fields outside of the object. The advantage of the method of retarded potentials is a possibility to avoid the problem of the boundary conditions.

For the case of a normal incidence of light at the quantum well surface, the method is described in⁷ and applied in¹². It was shown that the both versions of the semiclassical method lead to identical results. Calculations similar to the method of the retarded potentials are used in⁴.

Below, we consider the light scattering by a quantum dot of arbitrary form. We choose the electric field $\mathbf{E}_0(\mathbf{r}, t)$ in the form, allowing to obtain results for monochromatic and pulse irradiation.

II. CURRENT AND CHARGE DENSITIES INSIDE OF A QUANTUM DOT Плотности тока и заряда внутри КТ

According to⁷ (see Appendix 1), the average density of induced current inside of a quantum dot equals

$$\begin{aligned} \mathbf{j}(\mathbf{r}, t) &= \langle 0 | \mathbf{j}_1(\mathbf{r}, t) | 0 \rangle = \frac{ie^2}{2\pi\hbar\omega_g m_0^2} \sum_{\eta} F_{\eta}(\mathbf{r}) \\ &\times \int d^3\mathbf{r}' \int_{-\infty}^{\infty} dt' \int_{-\infty}^{\infty} d\omega e^{-i\omega(t-t')} \\ &\times \left\{ \frac{\mathbf{p}_{cv\eta}^* (\mathbf{p}_{cv\eta} \mathbf{E}_0(\mathbf{r}', t'))}{\omega - \omega_{\eta} + i\gamma_{\eta}/2} + \frac{\mathbf{p}_{cv\eta} (\mathbf{p}_{cv\eta}^* \mathbf{E}_0(\mathbf{r}', t'))}{\omega - \omega_{\eta} + i\gamma_{\eta}/2} \right\}, \quad (1) \end{aligned}$$

where $e(m_0)$ is the free electron charge (mass), $\hbar\omega_g$ is the energy gap, η is the exciton index, $F_{\eta}(\mathbf{r})$ is the real ("envelope") exciton wave function at $\mathbf{r}_e = \mathbf{r}_h = \mathbf{r}$, $\mathbf{r}_e(\mathbf{r}_h)$ is the electron (hole) radius-vector, $\mathbf{p}_{cv\eta}$ is the interband matrix element of the quasi-momentum operator, $\mathbf{E}(\mathbf{r}, t)$ is the electric field inside of a quantum dot; $\hbar\omega_{\eta}, \gamma_{\eta}$ is the energy (counted from the ground state energy) and exciton nonradiative damping, respectively.

Let us introduce the Fourier-transform $\mathcal{E}(\mathbf{r}, \omega)$ of the electric field¹⁷

$$\mathbf{E}(\mathbf{r}, t) = \frac{1}{2\pi} \int_{-\infty}^{\infty} d\omega e^{-i\omega t} \mathcal{E}(\mathbf{r}, \omega) + c.c.. \quad (2)$$

Then, we have from (1)

$$\begin{aligned} \mathbf{j}(\mathbf{r}, t) &= \frac{ie^2}{2\pi\hbar\omega_g m_0^2} \int_{-\infty}^{\infty} d\omega \sum_{\eta} F_{\eta}(\mathbf{r}) \\ &\times \left\{ \frac{\mathbf{p}_{cv\eta}^* \tilde{M}_{\eta}}{\omega - \omega_{\eta} + i\gamma_{\eta}/2} + \frac{\mathbf{p}_{cv\eta} \tilde{M}_{\eta}}{\omega + \omega_{\eta} + i\gamma_{\eta}/2} \right\}, \quad (3) \end{aligned}$$

where

$$\begin{aligned} M_{\eta} &= \int d^3r F_{\eta}(\mathbf{r}) (\mathbf{p}_{cv\eta} \mathcal{E}(\mathbf{r}, \omega)), \\ \tilde{M}_{\eta} &= \int d^3r F_{\eta}(\mathbf{r}) (\mathbf{p}_{cv\eta}^* \mathcal{E}(\mathbf{r}, \omega)). \quad (4) \end{aligned}$$

We determine the average density of induced charge inside of a quantum dot from the continuity equation

$$\text{div } \mathbf{j}(\mathbf{r}, t) + \frac{\partial \rho(\mathbf{r}, t)}{\partial t} = 0 \quad (5)$$

as

$$\begin{aligned} \rho(\mathbf{r}, t) &= \frac{e^2}{2\pi\hbar\omega_g m_0^2} \int_{-\infty}^{\infty} \frac{d\omega}{\omega} \\ &\times \sum_{\eta} \left\{ \text{div} (\mathbf{p}_{cv\eta}^* F_{\eta}(\mathbf{r})) \frac{M_{\eta}}{\omega - \omega_{\eta} + i\gamma_{\eta}/2} \right. \\ &\left. + \text{div} (\mathbf{p}_{cv\eta} F_{\eta}(\mathbf{r})) \frac{\tilde{M}_{\eta}}{\omega + \omega_{\eta} + i\gamma_{\eta}/2} \right\} + c.c. \quad (6) \end{aligned}$$

The genuine electric field inside of a quantum dot appears in the RHSs of expressions (1), (3) and (6). We assume that functions $F(\mathbf{r})$ and $dF(\mathbf{r})/d\mathbf{r}$ approach to 0 at $r \rightarrow \infty$, whence it follows that the current and charge densities equal 0 at $r \rightarrow \infty$.

III. RETARDED POTENTIALS

Vector and scalar potentials, corresponding to the induced electric field inside and outside of a quantum dot are defined by formulas¹³

$$\mathbf{A}(\mathbf{r}, t) = \frac{1}{c} \int d^3r' \frac{\mathbf{j}(\mathbf{r}', t - \nu|\mathbf{r} - \mathbf{r}'|/c)}{|\mathbf{r} - \mathbf{r}'|}, \quad (7)$$

$$\varphi(\mathbf{r}, t) = \frac{1}{\nu^2} \int d^3r' \frac{\rho(\mathbf{r}', t - \nu|\mathbf{r} - \mathbf{r}'|/c)}{|\mathbf{r} - \mathbf{r}'|}, \quad (8)$$

where ν is the refraction coefficient, which is identical inside and outside of a quantum dot. Substituting (3) and (6) in (7) and (8), we obtain

$$\begin{aligned} \mathbf{A}(\mathbf{r}, t) &= \frac{ie^2}{2\pi\hbar\omega_g m_0^2 c} \int_{-\infty}^{\infty} d\omega e^{-i\omega t} \\ &\times \sum_{\eta} \left\{ \frac{\mathbf{p}_{cv\eta}^* M_{\eta}}{\omega - \omega_{\eta} + i\gamma_{\eta}/2} + \frac{\mathbf{p}_{cv\eta} \tilde{M}_{\eta}}{\omega + \omega_{\eta} + i\gamma_{\eta}/2} \right\} \\ &\times \int d^3r' F_{\eta}(\mathbf{r}') \frac{e^{i\kappa|\mathbf{r} - \mathbf{r}'|}}{|\mathbf{r} - \mathbf{r}'|} + c.c., \quad (9) \end{aligned}$$

$$\begin{aligned} \varphi(\mathbf{r}, t) &= \frac{ie^2}{2\pi\hbar\omega_g m_0^2 \nu^2} \int_{-\infty}^{\infty} \frac{d\omega}{\omega} e^{-i\omega t} \\ &\times \sum_{\eta} \left\{ \frac{\mathbf{p}_{cv\eta}^* M_{\eta}}{\omega - \omega_{\eta} + i\gamma_{\eta}/2} + \frac{\mathbf{p}_{cv\eta} \tilde{M}_{\eta}}{\omega + \omega_{\eta} + i\gamma_{\eta}/2} \right\} \\ &\times \int d^3r' \frac{e^{i\kappa|\mathbf{r} - \mathbf{r}'|}}{|\mathbf{r} - \mathbf{r}'|} \frac{d}{d\mathbf{r}'} F_{\eta}(\mathbf{r}') + c.c., \quad (10) \end{aligned}$$

where $\kappa = \omega\nu/c$. In the RHS of (10) we transfer the derivative on $\mathbf{r}'$ with the opposite sign at the factor $e^{i\kappa|\mathbf{r}-\mathbf{r}'|}/|\mathbf{r}-\mathbf{r}'|$, and apply the ratio

$$\frac{d}{d\mathbf{r}'} \frac{e^{i\kappa|\mathbf{r}-\mathbf{r}'|}}{|\mathbf{r}-\mathbf{r}'|} = -\frac{d}{d\mathbf{r}} \frac{e^{i\kappa|\mathbf{r}-\mathbf{r}'|}}{|\mathbf{r}-\mathbf{r}'|}.$$

Получаем

$$\begin{aligned} \varphi(\mathbf{r}, t) &= \frac{ie^2}{2\pi\hbar\omega_g m_0^2 \nu^2} \int_{-\infty}^{\infty} \frac{d\omega}{\omega} e^{-i\omega t} \\ &\times \sum_{\eta} \left\{ \frac{\mathbf{p}_{cv\eta}^* M_{\eta}}{\omega - \omega_{\eta} + i\gamma_{\eta}/2} + \frac{\mathbf{p}_{cv\eta} \tilde{M}_{\eta}}{\omega + \omega_{\eta} + i\gamma_{\eta}/2} \right\} \\ &\times \frac{d}{d\mathbf{r}} \int d^3 r' \frac{e^{i\kappa|\mathbf{r}-\mathbf{r}'|}}{|\mathbf{r}-\mathbf{r}'|} F_{\eta}(\mathbf{r}') + c.c.. \end{aligned} \quad (11)$$

IV. ELECTRIC AND MAGNETIC FIELDS

Electric and magnetic fields are defined as

$$\begin{aligned} \mathbf{E}(\mathbf{r}, t) &= \frac{1}{c} \frac{\partial \mathbf{E}(\mathbf{r}, t)}{\partial t} - \frac{\partial \varphi(\mathbf{r}, t)}{\partial \mathbf{r}}, \\ \mathbf{H}(\mathbf{r}, t) &= \text{rot} \mathbf{A}. \end{aligned} \quad (12)$$

Let us calculate the fields $\mathbf{E}(\mathbf{r}, t)$ and $\mathbf{H}(\mathbf{r}, t)$ at very large distances r from a quantum dot, exceeding all the values of the length dimensionality, essential in our problem. It means that we determine fields in the limit $r \rightarrow \infty$.

Let us calculate the integral

$$J_{\eta}(\mathbf{r}) = \int d^3 r' F_{\eta}(\mathbf{r}') \frac{e^{i\kappa|\mathbf{r}-\mathbf{r}'|}}{|\mathbf{r}-\mathbf{r}'|}. \quad (13)$$

Since in the limit $r \rightarrow \infty$

$$e^{i\kappa|\mathbf{r}-\mathbf{r}'|} \simeq e^{i\kappa r - \kappa|\mathbf{r}-\mathbf{r}'|/r}$$

, we obtain the result

$$J_{\eta}(\mathbf{r})|_{r \rightarrow \infty} = \frac{e^{i\kappa r}}{r} P_{\eta}(\kappa_s), \quad (14)$$

where the designations

$$\begin{aligned} \kappa_s &= \kappa r / r, \\ P_{\eta}(\kappa) &= \int d^3 r F_{\eta}(\mathbf{r}) e^{-i\kappa r}, \end{aligned} \quad (15)$$

are introduced, and κ_s is the wave vector of the scattered light. We obtain also

$$\left(\frac{\partial}{\partial r} J_{\eta}(\mathbf{r}) \right)_{r \rightarrow \infty} = i\kappa \frac{\mathbf{r}}{r} \frac{e^{i\kappa r}}{r} P_{\eta}(\kappa_s), \quad (16)$$

and so on.

Using (14) and (16) and omitting contributions, small at $r \rightarrow \infty$, we obtain for the fields:

$$\begin{aligned} \mathbf{E}(\mathbf{r}, t) &= \frac{e^2}{2\pi\hbar\omega_g m_0^2 c^2} \int_{-\infty}^{\infty} d\omega \omega e^{-i\omega t} \frac{e^{i\kappa r}}{r} \sum_{\eta} P_{\eta}(\kappa_s) \\ &\times \left\{ \frac{[(\mathbf{p}_{cv}^* \frac{\mathbf{r}}{r}) \frac{\mathbf{r}}{r} - \mathbf{p}_{cv}^*] M_{\eta}}{\omega - \omega_{\eta} + i\gamma_{\eta}/2} + \frac{[(\mathbf{p}_{cv} \frac{\mathbf{r}}{r}) \frac{\mathbf{r}}{r} - \mathbf{p}_{cv}] \tilde{M}_{\eta}}{\omega + \omega_{\eta} + i\gamma_{\eta}/2} \right\} \\ &+ c.c., \end{aligned} \quad (17)$$

$$\begin{aligned} \mathbf{H}(\mathbf{r}, t) &= \frac{e^2 \nu}{2\pi\hbar\omega_g m_0^2 c^2} \int_{-\infty}^{\infty} d\omega \omega e^{-i\omega t} \frac{e^{i\kappa r}}{r} \sum_{\eta} P_{\eta}(\kappa_s) \\ &\times \left\{ \frac{[\mathbf{p}_{cv}^* \times \frac{\mathbf{r}}{r}] M_{\eta}}{\omega - \omega_{\eta} + i\gamma_{\eta}/2} + \frac{[\mathbf{p}_{cv} \times \frac{\mathbf{r}}{r}] \tilde{M}_{\eta}}{\omega + \omega_{\eta} + i\gamma_{\eta}/2} \right\} + c.c.. \end{aligned} \quad (18)$$

It follows from (17) and (18) that at $r \rightarrow \infty$ the fields $\mathbf{E}(\mathbf{r}, t)$ and $\mathbf{H}(\mathbf{r}, t)$ are perpendicular to the radius-vector $\mathbf{r}$.

V. DECOMPOSITION OF ELECTRIC AND MAGNETIC FIELDS ON THE COMPONENTS WITH LEFT AND RIGHT CIRCULAR POLARIZATIONS

Let us introduce the circular polarization vectors

$$\mathbf{e}_s^{\pm} = \frac{1}{\sqrt{2}} (\mathbf{e}_x \pm i\mathbf{e}_y)$$

, where the unite vectors $\mathbf{e}_x$ and $\mathbf{e}_y$ are perpendicular to the axis z , aligned parallel to $\mathbf{r}$.

Let us write down the fields as

$$\begin{aligned} \mathbf{E}(\mathbf{r}, t)_{r \rightarrow \infty} &= \mathbf{E}_c(\mathbf{r}, t) + \mathbf{E}_c^*(\mathbf{r}, t), \\ \mathbf{H}(\mathbf{r}, t)_{r \rightarrow \infty} &= \mathbf{H}_c(\mathbf{r}, t) + \mathbf{H}_c^*(\mathbf{r}, t), \end{aligned} \quad (19)$$

and decompose them on polarizations

$$\begin{aligned} \mathbf{E}_c(\mathbf{r}, t) &= E^+(\mathbf{r}, t) \mathbf{e}_s^+ + E^-(\mathbf{r}, t) \mathbf{e}_s^-, \\ \mathbf{H}_c(\mathbf{r}, t) &= H^+(\mathbf{r}, t) \mathbf{e}_s^+ + H^-(\mathbf{r}, t) \mathbf{e}_s^-. \end{aligned} \quad (20)$$

Multiplying both parts of (20) consecutively on $\mathbf{e}_s^+$ and $\mathbf{e}_s^-$, we find

$$\begin{aligned} \mathbf{E}^{\pm}(\mathbf{r}, t) &= -\frac{e^2}{2\pi\hbar\omega_g m_0^2 c^2 r} \int_{-\infty}^{\infty} d\omega \omega e^{i\kappa r - i\omega t} \\ &\times Q(\kappa_s, \omega, \mathbf{e}_s^{\pm}), \end{aligned} \quad (21)$$

$$H^{\pm}(\mathbf{r}, t) = \mp i\nu E^{\pm}(\mathbf{r}, t), \quad (22)$$

where

$$\begin{aligned} Q(\kappa_s, \omega, \mathbf{e}_s^{\pm}) &= \sum_{\eta} P_{\eta}(\kappa_s) \\ &\times \left\{ \frac{(\mathbf{p}_{cv\eta} \mathbf{e}_s)^* M_{\eta}}{\omega - \omega_{\eta} + i\gamma_{\eta}/2} + \frac{(\mathbf{p}_{cv\eta} \mathbf{e}_s^{\pm*}) \tilde{M}_{\eta}}{\omega + \omega_{\eta} + i\gamma_{\eta}/2} \right\} + c.c.. \end{aligned} \quad (23)$$

VI. THE POINTING VECTOR

The Pointing vector is equal¹³

$$\mathbf{S}(\mathbf{r}, t) = \frac{c}{4\pi} \mathbf{E}(\mathbf{r}, t) \times \mathbf{H}(\mathbf{r}, t). \quad (24)$$

Substituting (19) in (24), we obtain

$$\begin{aligned} \mathbf{S}(\mathbf{r}, t)_{r \rightarrow \infty} &= \frac{c}{4\pi} \\ &\times \{ \mathbf{E}_c(\mathbf{r}, t) \times \mathbf{H}_c^*(\mathbf{r}, t) + \mathbf{E}_c^*(\mathbf{r}, t) \times \mathbf{H}_c(\mathbf{r}, t) \\ &+ \mathbf{E}_c(\mathbf{r}, t) \times \mathbf{H}_c(\mathbf{r}, t) + \mathbf{E}_c^*(\mathbf{r}, t) \times \mathbf{H}_c^*(\mathbf{r}, t) \}. \end{aligned} \quad (25)$$

Two first terms in the RHS of (25) provide the main contribution into the Pointing vector, averaged on time.¹⁸ Restricting by that contribution, we have

$$\mathbf{S}(\mathbf{r}, t)_{r \rightarrow \infty} = \sum_{\mu} \mathbf{S}(\mathbf{r}, t, \mathbf{l}_s), \quad (26)$$

where $\sum_{\mu}$ designates summation on circular polarizations of scattered light,

$$\begin{aligned} \mathbf{S}(\mathbf{r}, t, \mathbf{l}_s)_{r \rightarrow \infty} &= \frac{c}{(2\pi)^3} \left(\frac{e^2}{\hbar c} \right)^2 \frac{\nu}{m_g m_0^4 c} \frac{\mathbf{r}}{r} \\ &\times \left| \int_{-\infty}^{\infty} d\omega \omega e^{i\kappa r - i\omega t} Q(\kappa_s, \omega, \mathbf{e}_s) \right|^2. \end{aligned} \quad (27)$$

Opening the square of module in the RHS of (27), we obtain

$$\begin{aligned} &\int_{-\infty}^{\infty} d\omega \int_{-\infty}^{\infty} d\omega' \omega \omega' e^{i(\kappa - \kappa')r - i(\omega - \omega')t} \\ &\times Q(\kappa_s, \omega, \mathbf{e}_s) Q^*(\kappa'_s, \omega', \mathbf{e}_s), \end{aligned} \quad (28)$$

where $\kappa' = \omega' \nu / c$, $\kappa'_s = \kappa'_s \mathbf{r} / r$. As it will be seen below, the value $Q(\kappa_s, \omega, \mathbf{e}_s)$ always contains the factor $D_0(\omega)$, describing the form of the stimulating pulse $\mathbf{E}_0(\mathbf{r}, t)$, defined as

$$\mathbf{E}_0(\mathbf{r}, t) = E_0 \mathbf{e}_{\ell} \int_{-\infty}^{\infty} d\omega e^{i\kappa_{\ell} \mathbf{r} - i\omega t} D_0(\omega) + c.c., \quad (29)$$

where κ_{ℓ} is the vector, which magnitude is equal $\omega_{\ell} \nu / c$, and the direction is defined by the direction of stimulating light. For the monochromatic irradiation

$$D_0(\omega) = \delta(\omega - \omega_{\ell}), \quad (30)$$

and for the pulse irradiation the frequency is spreaded beside ω_{ℓ} for some interval $\Delta\omega$, inversely proportional to the pulse duration Δt . In any case, the carrying frequency ω_{ℓ} disappears from (28) due to the factor $e^{-i(\omega - \omega')t}$ in the integrand. The contribution of the last two terms from the RHS of (25) contains under integral $e^{-i(\omega + \omega')t}$, what makes this contribution rapidly oscillating in time and approaching 0 at averaging on time.

VII. THE SCATTERING CROSS SECTION

In a case of monochromatic irradiation, when condition (30) is satisfied, it is conveniently to introduce a concept of the cross section of light scattering on a quantum dot.

The differential cross section is defined as a ratio of a magnitude of scattered energy flux into an solid angle interval do_s to the magnitude of incident energy flux on the area unite in a time unite. For a monochromatic irradiation, it is equal

$$S_0 = \frac{c\nu}{2\pi} E_0^2. \quad (31)$$

Using (26) and (27), we obtain

$$\begin{aligned} \frac{d\sigma}{do_s} &= \sum_{\mu} \frac{d\sigma_{\mu}}{do_s}, \\ \frac{d\sigma_{\mu}}{do_s} &= \frac{1}{(2\pi)^2} \left(\frac{e^2}{\hbar c} \right)^2 \frac{1}{\omega_g^2 m_0^4 c^2 E_0^2} \\ &\times \left| \int_{-\infty}^{\infty} d\omega \omega e^{i(\kappa r - i\omega t)} Q(\kappa_s, \omega, \mathbf{e}_s) \right|^2. \end{aligned} \quad (32)$$

In a case of monochromatic irradiation, the factor $\delta(\omega - \omega_{\ell})$ presents always in the expression for $Q(\kappa_s, \omega', \mathbf{e}_s)$ (see below). It means that we can write

$$Q(\kappa_s, \omega, \mathbf{e}_s) = q(\kappa_s, \omega_{\ell}, \mathbf{e}_s) \delta(\omega - \omega_{\ell}), \quad (33)$$

and then (32) is transformed to the form

$$\frac{d\sigma_{\mu}}{do_s} = \frac{1}{(2\pi)^2} \left(\frac{e^2}{\hbar c} \right)^2 \frac{\omega_{\ell}^2}{\omega_g^2 m_0^4 c^2 E_0^2} |q(\kappa_s, \omega_{\ell}, \mathbf{e}_s)|^2, \quad (34)$$

and dependencies on r and t disappear.

VIII. THE LOWEST APPROXIMATION ON THE ELECTRON-LIGHT INTERACTION

In the lowest approximation the genuine electric field inside of a quantum dot, which is included in expressions for $M_{\eta}(\omega)$ and $\tilde{M}_{\eta}(\omega)$, is substituted by the stimulating field. According to (29) and (2), the Fourier transform is as follows

$$\mathcal{E}_0(\mathbf{r}, \omega,) = 2\pi \mathbf{e}_{\ell} E_0 e^{i\kappa_{\ell} \mathbf{r}} D_0(\omega). \quad (35)$$

Then,

$$\begin{aligned} M_{\eta 0} &= 2\pi E_0 D_0(\omega) P_{\eta}^*(\kappa_{\ell}) (\mathbf{e}_{\ell} \mathbf{p}_{cv\eta}), \\ \tilde{M}_{\eta 0} &= 2\pi E_0 D_0(\omega) P_{\eta}^*(\kappa_{\ell}) (\mathbf{e}_{\ell} \mathbf{p}_{cv\eta}^*), \end{aligned} \quad (36)$$

$$\begin{aligned} Q_0 &= 2\pi E_0 D_0(\omega) \sum_{\eta} P_{\eta}(\kappa_s) P_{\eta}^*(\kappa_{\ell}) \\ &\times \left\{ \frac{(\mathbf{e}_s \mathbf{p}_{cv\eta}^*)(\mathbf{e}_{\ell} \mathbf{p}_{cv\eta})}{\omega - \omega_{\eta} + i\gamma_{\eta}/2} + \frac{(\mathbf{e}_s^* \mathbf{p}_{cv\eta})(\mathbf{e}_{\ell} \mathbf{p}_{cv\eta}^*)}{\omega + \omega_{\eta} + i\gamma_{\eta}/2} \right\}. \end{aligned} \quad (37)$$

Substituting (37) in (21), (22) and (27), we obtain electric and magnetic fields and the Poynting vector in the limit $r \rightarrow \infty$.

For the monochromatic irradiation, we obtain the differential section

$$\frac{d\sigma_{\mu 0}}{do_s} = \left(\frac{e^2}{\hbar c} \right)^2 \frac{\omega_\ell^2}{\omega_g^2 m_0^4 c^2} \left| \sum_\eta P_\eta(\kappa_s) P_\eta^*(\kappa_\ell) (\mathbf{v}_\eta \mathbf{e}_s^*) \right|^2, \quad (38)$$

where

$$|\kappa_s| = |\kappa_\ell| = \frac{\omega_\ell \nu}{c},$$

$$\mathbf{v}_\eta = \frac{\mathbf{p}_{cv\eta}^* (\mathbf{e}_\ell \mathbf{p}_{cv\eta})}{\omega - \omega_\eta + i\gamma_\eta/2} + \frac{\mathbf{p}_{cv\eta} (\mathbf{e}_\ell \mathbf{p}_{cv\eta}^*)}{\omega + \omega_\eta + i\gamma_\eta/2}. \quad (39)$$

IX. COMPARISON OF RESULTS OF SEMICLASSICAL AND QUANTUM THEORIES¹⁴

Let us omit in (38) the nonresonant term, containing $(\omega + \omega_\eta + i\gamma_\eta/2)^{-1}$. We obtain

$$\frac{d\sigma_{\mu 0}}{do_s} = \left(\frac{e^2}{\hbar c} \right)^2 \frac{\omega_\ell^2}{\omega_g^2 m_0^4 c^2} \times \left| \sum_\eta \frac{P_\eta(\kappa_s) P_\eta^*(\kappa_\ell) (\mathbf{p}_{cv\eta} \mathbf{e}_\ell) (\mathbf{p}_{cv\eta} \mathbf{e}_s)^*}{\omega - \omega_\eta + i\gamma_\eta/2} \right|^2. \quad (40)$$

It follows from (40) that if the light wave length exceeds the quantum dot sizes and

$$P_\eta(\kappa_\ell) \simeq P_\eta(\kappa_s) \simeq P_\eta(0),$$

the polarization and angular distribution of scattered light depend only on vectors $\mathbf{p}_{cv\eta}$, corresponding to the exciton, which is in resonance with the stimulating light. This result is very close to results of the quantum theory (see (15) from¹⁵), but the factor $(\omega_\ell/\omega_g)^4$ is substituted by $(\omega_\ell/\omega_g)^2$, what is incidentally under resonant conditions, and the value $(\omega - \omega_\eta + i\delta)^{-1}$ is substituted by $(\omega - \omega_\eta + i\gamma_\eta/2)^{-1}$. Obviously, the semiclassical method provides more exact results even in the lowest approximation on the electron-light interaction, since it allows to include into the theory the nonradiative damping γ_η (which determines light absorption by a quantum dot). Besides, Kpome toro, the semiclassical method allows to investigate light scattering and absorption for a pulse irradiation.

Further, we pass on to the precise semiclassical theory, taking into account all the orders on the electron-light interaction.

X. THE EQUATION FOR ELECTRIC FIELD INSIDE OF A QUANTUM DOT

First of all, we calculate defined in (4) M_η and $\tilde{M}_\eta$ from the RHS of (23), i.e., we substitute in (4)

the Fourier-components of the genuine electric field inside of the quantum dot. We obtain an equation for these Fourier-components $\mathcal{E}(\mathbf{r}, \omega)$ as follows, writing the genuine electric field inside of the quantum dot in the form

$$\mathbf{E}(\mathbf{r}, t) = \mathbf{E}_0(\mathbf{r}, t) - \frac{1}{c} \frac{\partial \mathbf{A}(\mathbf{r}, t)}{\partial t} - \frac{\partial \varphi(\mathbf{r}, t)}{\partial \mathbf{r}}, \quad (41)$$

where $\mathbf{E}_0(\mathbf{r}, t)$ is the stimulating electric field (29), $\mathbf{A}(\mathbf{r}, t)$ and $\varphi(\mathbf{r}, t)$ are the vector and scalar potentials, defined in (9) and (10) and corresponding to the secondary radiation of the quantum dot. For the Fourier-components $\mathcal{E}(\mathbf{r}, \omega)$ of the field (41), we obtain the integral equation

$$\mathcal{E}(\mathbf{r}, \omega) = 2\pi \mathbf{e}_\ell E_0 e^{i\kappa_\ell \mathbf{r}} D_0(\omega) - \frac{e^2}{\hbar c} \frac{\omega}{\omega_g m_0^2 c} \times \sum_\eta \left\{ \frac{M_\eta \left[\mathbf{p}_{cv\eta}^* + \kappa^{-2} \frac{\partial}{\partial \mathbf{r}} \left(\frac{\partial \mathbf{p}_{cv\eta}^*}{\partial \mathbf{r}} \right) \right]}{\omega - \omega_\eta + i\gamma_\eta/2} + \frac{\tilde{M}_\eta \left[\mathbf{p}_{cv\eta} + \kappa^{-2} \frac{\partial}{\partial \mathbf{r}} \left(\frac{\partial \mathbf{p}_{cv\eta}}{\partial \mathbf{r}} \right) \right]}{\omega + \omega_\eta + i\gamma_\eta/2} \right\} \Phi_\eta(\mathbf{r}), \quad (42)$$

where

$$\Phi_\eta(\mathbf{r}) = \int d^3 r' F_\eta(\mathbf{r}') \frac{e^{i\kappa|\mathbf{r}-\mathbf{r}'|}}{|\mathbf{r}-\mathbf{r}'|}. \quad (43)$$

XI. THE EQUATION SYSTEM FOR COEFFICIENTS M_η AND $\tilde{M}_\eta$

We multiply (42) at $F_{\eta'}(\mathbf{r}) \mathbf{p}_{cv\eta'}$ and integrate on $\mathbf{r}$, then, perform the same operation, using $F_{\eta'}(\mathbf{r}) \mathbf{p}_{cv\eta'}^*$. We obtain the following system of equations for M_η and $\tilde{M}_\eta$ ¹⁹

$$M_{\eta'} = 2\pi E_0 D_0(\omega) (\mathbf{e}_\ell \mathbf{p}_{cv\eta'}) P_{\eta'}^*(\kappa_\ell) - \frac{e^2}{\hbar c} \frac{\omega}{\omega_g m_0^2 c} \times \sum_\eta \left\{ \frac{M_\eta R_{\eta\eta'}}{\omega - \omega_\eta + i\gamma_\eta/2} + \frac{\tilde{M}_\eta S_{\eta\eta'}}{\omega + \omega_\eta + i\gamma_\eta/2} \right\},$$

$$\tilde{M}_{\eta'} = 2\pi E_0 D_0(\omega) (\mathbf{e}_\ell \mathbf{p}_{cv\eta'}^*) P_{\eta'}^*(\kappa_\ell) - \frac{e^2}{\hbar c} \frac{\omega}{\omega_g m_0^2 c} \times \sum_\eta \left\{ \frac{M_\eta \tilde{S}_{\eta\eta'}}{\omega - \omega_\eta + i\gamma_\eta/2} + \frac{\tilde{M}_\eta \tilde{R}_{\eta\eta'}}{\omega + \omega_\eta + i\gamma_\eta/2} \right\}, \quad (44)$$

where

$$R_{\eta\eta'} = (\mathbf{p}_{cv\eta}^* \mathbf{p}_{cv\eta'}) \int d^3 r F_{\eta'}(\mathbf{r}) \Phi_\eta(\mathbf{r}) + \frac{1}{\kappa^2} \int d^3 r F_{\eta'}(\mathbf{r}) \left(\frac{\partial \mathbf{p}_{cv\eta'}}{\partial \mathbf{r}} \right) \left(\frac{\partial \mathbf{p}_{cv\eta}}{\partial \mathbf{r}} \right) \Phi_\eta(\mathbf{r}),$$

$$S_{\eta\eta'} = (\mathbf{p}_{cv\eta} \mathbf{p}_{cv\eta'}) \int d^3 r F_{\eta'}(\mathbf{r}) \Phi_\eta(\mathbf{r}) + \frac{1}{\kappa^2} \int d^3 r F_{\eta'}(\mathbf{r}) \left(\frac{\partial \mathbf{p}_{cv\eta'}}{\partial \mathbf{r}} \right) \left(\frac{\partial \mathbf{p}_{cv\eta}}{\partial \mathbf{r}} \right) \Phi_\eta(\mathbf{r}). \quad (45)$$

The integrals on $\mathbf{r}$ from (45) may be written in a more symmetrical form

$$\begin{aligned} \int d^3r F_{\eta'}(\mathbf{r}) \Phi_{\eta}(\mathbf{r}) &= \int d^3r \int d^3r' F_{\eta'}(\mathbf{r}) F_{\eta}(\mathbf{r}) \\ &\times \frac{e^{i\kappa|\mathbf{r}-\mathbf{r}'|}}{|\mathbf{r}-\mathbf{r}'|}, \\ \int d^3r F_{\eta'}(\mathbf{r}) \left(\frac{\partial \mathbf{p}_{cv\eta}}{\partial \mathbf{r}} \right) \left(\frac{\partial \mathbf{p}_{cv\eta}^*}{\partial \mathbf{r}} \right) \Phi_{\eta}(\mathbf{r}) \\ &= \int d^3r \int d^3r' F_{\eta'}(\mathbf{r}) F_{\eta}(\mathbf{r}) \left(\mathbf{p}_{cv\eta'} \frac{\partial}{\partial \mathbf{r}} \right) \left(\mathbf{p}_{cv\eta'} \frac{\partial}{\partial \mathbf{r}} \right) \\ &\times \frac{e^{i\kappa|\mathbf{r}-\mathbf{r}'|}}{|\mathbf{r}-\mathbf{r}'|}. \end{aligned} \quad (46)$$

The coefficients from (44) have properties

$$R_{\eta'\eta} = \tilde{R}_{\eta\eta'}, S_{\eta'\eta} = S_{\eta\eta'}. \quad (47)$$

XII. CALCULATION OF THE FUNCTION $\Phi(\mathbf{r})$

Let us obtain the Fourier-transformation of the function

$$\Phi(\mathbf{r}) = \int d^3r' F(\mathbf{r}') \frac{e^{i\kappa|\mathbf{r}-\mathbf{r}'|}}{|\mathbf{r}-\mathbf{r}'|}.$$

We obtain

$$\Phi(\mathbf{r}) = \frac{1}{(2\pi)^3} \int d^3\kappa F(\kappa) f_{\kappa}(\kappa) e^{i\kappa\mathbf{r}}, \quad (48)$$

where

$$\begin{aligned} F_{\kappa}(\kappa) &= \int d^3r e^{-i\kappa\mathbf{r}} F(\mathbf{r}) = P(\kappa), f_{\kappa}(\kappa) \\ &= \int d^3r e^{-i\kappa\mathbf{r}} \frac{e^{i\kappa\mathbf{r}}}{r}. \end{aligned}$$

Integration on $\mathbf{r}$ results in

$$\begin{aligned} f_{\kappa}(\mathbf{k}) &= f_{\kappa}(k) \\ &= \frac{2\pi}{k} \left\{ \frac{P}{k-\kappa} + \frac{P}{k+\kappa} + i\pi\delta(k-\kappa) - \frac{P}{k+\kappa} \right\}, \\ \frac{P}{k-\kappa} &= \frac{1}{2} \left(\frac{1}{k-\kappa+i\delta} + \frac{P}{k-\kappa-i\delta} \right), \\ \delta &\rightarrow +0. \end{aligned} \quad (49)$$

The last term in the RHS of (49) may be omitted, then

$$\begin{aligned} \Phi(\mathbf{r}) &= \frac{1}{(2\pi)^3} \int d^3k e^{i\mathbf{k}\mathbf{r}} P(\mathbf{k}) \\ &\times \left\{ 4\pi \frac{P}{k^2-\kappa^2} + \frac{2i\pi^2}{\kappa} \delta(k-\kappa) \right\}. \end{aligned} \quad (50)$$

It is easy to check that the function $\Phi(\mathbf{r})$ satisfies the equation

$$(\kappa^2 + \Delta)\Phi(\mathbf{r}) = 4\pi F(\mathbf{r}), \Delta = \left(\frac{\partial}{\partial \mathbf{r}} \right)^2. \quad (51)$$

XIII. THE EQUATION SYSTEM FOR ONE DEGENERATED ENERGY LEVEL

Let us consider n times degenerated exciton energy level. Without taking into account the electron-light interaction,

$$\omega_{\eta} = \omega_0, \quad (52)$$

and

$$\gamma_{\eta} = \gamma, F_{\eta}(\mathbf{r}) = F(\mathbf{r}), P_{\eta}(\kappa) = P(\kappa), \Phi_{\eta}(\mathbf{r}) = \Phi(\mathbf{r}). \quad (53)$$

Only the vectors $\mathbf{p}_{cv\eta}$ depend on index η , taking n values. Taking into account only this level (assuming, that the frequency of stimulating light is in the resonance with ω_0), we obtain from (44) and (50) 2n algebraical equations for 2n quantities M_{η} and $\tilde{M}_{\eta}$.

$$\begin{aligned} M_{\eta'} &= 2\pi E_0 D_0(\omega) (\mathbf{l}_{\ell} \mathbf{p}_{cv\eta'}) P^*(\kappa_{\ell}) \\ &+ \sum_{\eta} \left\{ \frac{M_{\eta} \Omega(\mathbf{p}_{cv\eta}^*, \mathbf{p}_{cv\eta'})}{\omega - \omega_0 + i\gamma/2} + \frac{\tilde{M}_{\eta} \Omega(\mathbf{p}_{cv\eta}, \mathbf{p}_{cv\eta'})}{\omega + \omega_0 + i\gamma/2} \right\}, \\ \tilde{M}_{\eta'} &= 2\pi E_0 D_0(\omega) (\mathbf{e}_{\ell} \mathbf{p}_{cv\eta'}) P^*(\kappa_{\ell}) \\ &+ \sum_{\eta} \left\{ \frac{M_{\eta} \Omega(\mathbf{p}_{cv\eta}^*, \mathbf{p}_{cv\eta'})}{\omega - \omega_0 + i\gamma/2} + \frac{\tilde{M}_{\eta} \Omega(\mathbf{p}_{cv\eta}, \mathbf{p}_{cv\eta'})}{\omega + \omega_0 + i\gamma/2} \right\}, \end{aligned} \quad (54)$$

where

$$\Omega(\mathbf{p}_1, \mathbf{p}_2) = \Omega(\mathbf{p}_2, \mathbf{p}_1) = \Delta\omega(\mathbf{p}_1, \mathbf{p}_2) - i\gamma_r(\mathbf{p}_1, \mathbf{p}_2)/2, \quad (55)$$

$$\begin{aligned} \Delta\omega(\mathbf{p}_1, \mathbf{p}_2) &= \frac{e^2}{2\pi^2 \hbar c} \frac{\omega}{\omega_g m_0^2 c \kappa^2} \int d^3k |P(\mathbf{k})|^2 \\ &\times \left\{ (\mathbf{p}_1, \mathbf{p}_2) + [(\mathbf{k}_1 \mathbf{p}_1)(\mathbf{k} \mathbf{p}_2) - k^2(\mathbf{p}_1, \mathbf{p}_2)] \frac{P}{\kappa^2 - k^2} \right\}, \end{aligned} \quad (56)$$

$$\begin{aligned} \gamma_r(\mathbf{p}_1, \mathbf{p}_2) &= \frac{e^2}{2\pi \hbar c} \frac{\omega}{\omega_g m_0^2 c \kappa} \int d\omega_{\kappa} |P(\mathbf{k})|^2 \\ &\times \{ \kappa^2(\mathbf{p}_1 \mathbf{p}_2) - (\kappa \mathbf{p}_1)(\kappa \mathbf{p}_2) \}. \end{aligned} \quad (57)$$

XIV. THE EXCITON $\Gamma_6 \times \Gamma_7$ IN CUBIC CRYSTALS OF T_d CLASS

As an example, let us consider an exciton, formed by an electron from twice degenerated Γ_6 conduction band and by a hole from twice degenerated Γ_7 valence band, splitted by the spin - orbital interaction.

The electron wave functions have the structure?

$$\Psi_{c1} = iS \uparrow, \Psi_{c2} = iS \downarrow, \quad (58)$$

and the the hole wave functions are

$$\begin{aligned} \Psi_{h1} &= \frac{1}{\sqrt{3}} (X - iY) \uparrow - \frac{1}{\sqrt{3}} Z \downarrow, \\ \Psi_{h2} &= \frac{1}{\sqrt{3}} (X + iY) \downarrow + \frac{1}{\sqrt{3}} Z \uparrow. \end{aligned} \quad (59)$$

Combining (58) и (59) in pairs, we obtain four times degenerated excitonic state, for which the vectors $\mathbf{p}_{cv\eta}$ are

$$\begin{aligned}\mathbf{p}_{cv1} &= \frac{p_{cv}}{\sqrt{3}}(\mathbf{e}_x - i\mathbf{e}_y), \\ \mathbf{p}_{cv2} &= \frac{p_{cv}}{\sqrt{3}}(\mathbf{e}_x + i\mathbf{e}_y), \\ \mathbf{p}_{cv3} &= \frac{p_{cv}}{\sqrt{3}}\mathbf{e}_z, \\ \mathbf{p}_{cv4} &= -\frac{p_{cv}}{\sqrt{3}}\mathbf{e}_z,\end{aligned}\quad (60)$$

where we introduced the scalar

$$p_{cv} = i\langle S|\hat{p}_x|X\rangle, \quad (61)$$

$\mathbf{e}_x, \mathbf{e}_y, \mathbf{e}_z$ are the unite vectors along the crystallographic axes.

We consider the circular polarization of stimulating and scattered light, i.e.,

$$\begin{aligned}\mathbf{e}_\ell^\pm &= \frac{1}{\sqrt{2}}(\mathbf{e}_{x\ell} \pm i\mathbf{e}_{y\ell}), \\ \mathbf{e}_s^\pm &= \frac{1}{\sqrt{2}}(\mathbf{e}_{xs} \pm i\mathbf{e}_{ys}),\end{aligned}\quad (62)$$

where the unite vectors $\mathbf{e}_{x\ell}$ and $\mathbf{e}_{y\ell}$ are perpendicular to the axis z_ℓ along the vector κ_ℓ , the unite vectors $\mathbf{e}_{xs}$ and $\mathbf{e}_{ys}$ are perpendicular to the axis z_s along the vector κ_s .

In the case of the exciton $\Gamma_6 \times \Gamma_7$, it is conveniently to sum in (3) and (6) on indexes η , using (55). Calculations, described in sections III - XIII, result in the same expressions, certainly. After summation, we have

$$\begin{aligned}\mathbf{j}(\mathbf{r}, t) &= \frac{ie^2 p_{cv}^2}{3\pi\hbar\omega_g m_0^2} F(\mathbf{r}) \int_{-\infty}^{\infty} d\omega e^{-i\omega t} \mathbf{T}(\omega) L(\omega) \\ &+ c.c.,\end{aligned}\quad (63)$$

$$\begin{aligned}\rho(\mathbf{r}, t) &= \frac{ie^2 p_{cv}^2}{3\pi\hbar\omega_g m_0^2} F(\mathbf{r}) \int_{-\infty}^{\infty} \frac{d\omega}{\omega} \left(\frac{dF(\mathbf{r})}{d\mathbf{r}} \right) \mathbf{T}(\omega) L(\omega) \\ &+ c.c.,\end{aligned}\quad (64)$$

where the designations are introduced

$$\begin{aligned}\mathbf{T}(\omega) &= \int d^3r \mathcal{E}(\mathbf{r}, \omega) F(\mathbf{r}), \\ L(\omega) &= \frac{1}{\omega - \omega_0 + i\gamma/2} + \frac{1}{\omega + \omega_0 + i\gamma/2}.\end{aligned}\quad (65)$$

Expressions (63) and (64) do not contain evidently the vectors $\mathbf{r}_{cv\eta}$ and allow to calculate the light scattering cross section by a quantum dot in the resonance with the exciton $\Gamma_6 \times \Gamma_7$. It follows from (63) and (64), that in that case all the values do not depend on the crystallographic axes direction.

Using (7) and (8), we obtain vector and scalar potentials

$$\begin{aligned}\mathbf{A}(\mathbf{r}, t) &= \frac{ie^2 p_{cv}^2}{3\pi\hbar\omega_g m_0^2 c} \int_{-\infty}^{\infty} d\omega e^{-i\omega t} L(\omega) \mathbf{T}(\omega) \\ &\times \int d^3r' F(\mathbf{r}') \frac{e^{i\kappa|\mathbf{r}-\mathbf{r}'|}}{|\mathbf{r}-\mathbf{r}'|} + c.c.,\end{aligned}\quad (66)$$

$$\begin{aligned}\varphi(\mathbf{r}, t) &= \frac{ie^2 p_{cv}^2}{3\pi\hbar\omega_g m_0^2 \nu^2} \int_{-\infty}^{\infty} \frac{d\omega}{\omega} e^{-i\omega t} L(\omega) \mathbf{T}(\omega) \\ &\times \frac{d}{d\mathbf{r}} \int d^3r' F(\mathbf{r}') \frac{e^{i\kappa|\mathbf{r}-\mathbf{r}'|}}{|\mathbf{r}-\mathbf{r}'|} + c.c.. \end{aligned}\quad (67)$$

For electric and magnetic fields, one can apply (19) - (22) with the substitution

$$Q(\kappa_s, \omega, \mathbf{e}_s^\pm) = \frac{2p_{cv}^2}{3} P(\kappa_s) (\mathbf{T}(\omega) \mathbf{e}_s) L(\omega). \quad (68)$$

For the Pointing vector at $r \rightarrow \infty$, one has to use (27) with substitution (68). Для сечения рассеяния при монохроматическом облучении применимы точные формулы Formulas (33) - (34) with substitution (68) are applicable for the scattering section at a monochromatic irradiation. In the lowest order on the electron-light interaction, we obtain

$$\begin{aligned}\frac{d\sigma_\mu}{d\omega_s} &= \frac{4}{9} \left(\frac{e^2}{\hbar c} \right)^2 \frac{p_{cv}^4 \omega_\ell^2}{\omega_g^2 m_0^4 c^2} |\mathbf{e}_\ell \mathbf{e}_s^*|^2 |P(\kappa_\ell)|^2 |P(\kappa_s)|^2 \\ &\times \left| \frac{1}{\omega - \omega_0 + i\gamma/2} + \frac{1}{\omega + \omega_0 + i\gamma/2} \right|^2,\end{aligned}\quad (69)$$

and

$$\begin{aligned}|\mathbf{e}_\ell^+ \mathbf{e}_s^-|^2 &= \frac{1}{4} (1 + \cos \theta)^2, \\ |\mathbf{e}_\ell^+ \mathbf{e}_s^+|^2 &= \frac{1}{4} (1 - \cos \theta)^2,\end{aligned}\quad (69a)$$

where θ is the scattering angle, what coincide with results (32) - (35) of the article¹⁴, if in (69) the nonresonant contribution (proportional to $(\omega + \omega_0 + i\gamma/2)^{-1}$) is omitted, and the factor $(\omega_\ell/\omega_g)^2$ is substituted by $(\omega_\ell/\omega_g)^4$, and $(\omega - \omega_0 + i\gamma/2)^{-1}$ is substituted by $(\omega + \omega_0 + i\delta)^{-1}$, $\delta \rightarrow 0$.

XV. THE EQUATION FOR THE ELECTRIC FIELD INSIDE OF A QUANTUM DOT IN THE CASE OF THE EXCITON $\Gamma_6 \times \Gamma_7$

Having substituted expressions (29), (66) and (67) in (41), we obtain the equation for the Fourier-component of the electric field

Подставив в (41) выражения (29), (66) и (67), получим уравнение для Фурье-компонент электрического поля

$$\mathcal{E}(\mathbf{r}, \omega) = 2\pi E_0 \mathbf{e}_\ell e^{i\kappa_\ell \mathbf{r}} D_0(\omega) - \frac{2}{3} \frac{e^2}{\hbar c} \frac{p_{cv}^2 \omega}{m_0^2 \omega_g c} L(\omega) \times \left[\mathbf{T}(\omega) + \kappa^{-2} \left(\frac{d}{d\mathbf{r}} \mathbf{T}(\omega) \right) \right] \Phi(\mathbf{r}). \quad (70)$$

The equation (70) can be obtained also from (42) with the help of substitution the vector $\mathbf{p}_{cv\eta}$ from (60) and summation on indexes η , taking the values from 1 to 4.

Let us multiply (70) on $F(\mathbf{r})$ and integrate on $\mathbf{r}$. We obtain the equation for the vector $\mathbf{T}(\omega)$

$$\mathbf{T}(\omega) = 2\pi E_0 \mathbf{e}_\ell D_0 P^*(\mathbf{r}_\ell) - C(\omega) \int d^3 r \left[\mathbf{T}(\omega) \Phi(\mathbf{r}) + \kappa^{-2} \left(\frac{d}{d\mathbf{r}} \mathbf{T}(\omega) \right) \Phi(\mathbf{r}) \right], \quad (71)$$

where the designation

$$C(\omega) = \frac{2}{3} \left(\frac{e^2}{\hbar c} \right) \frac{p_{cv}^2 \omega}{\omega_g^2 m_0^2 c} L(\omega), \quad (72)$$

is introduced, and expression (50) is used for the function $\Phi(\mathbf{r})$. Substituting (50) in (71), we obtain

$$\mathbf{T}(\omega)(1 + C(\omega)) \int d^3 k J(\mathbf{k}) = 2\pi E_0 \mathbf{e}_\ell D_0 P^*(\kappa_\ell) + \frac{C(\omega)}{\kappa^2} \int d^3 k \mathbf{k} (\mathbf{k} \mathbf{T}(\omega)) J(\mathbf{k}), \quad (73)$$

$$J(\mathbf{k}) = |P(\mathbf{k})|^2 \left[\frac{1}{2\pi^2} \frac{P}{k^2 - \kappa^2} + \frac{i}{4\pi\kappa} \delta(k - \kappa) \right]. \quad (74)$$

Expression (73) may be treated as a system of three equations for three components $T_x(\omega), T_y(\omega), T_z(\omega)$ of vector $\mathbf{T}(\omega)$. The result must be substituted into the formula for the scattering section at monochromatic excitation or into the formula for the Pointing vector at $r \rightarrow \infty$ at pulse excitation. Thus, thus the problem of the resonant light scattering is solved.

XVI. LIGHT SCATTERING BY THE EXCITON $\Gamma_6 \times \Gamma_7$ IN A SPHERICAL QUANTUM DOT

Solution of the problem is simplified, when $P(\mathbf{k})$ depends only on the magnitude of the vector $\mathbf{k}$ in the case of a spherically symmetrical function

$$F(\mathbf{r}) = F(r). \quad (75)$$

Then, at $J(\mathbf{k}) = J(k)$,

$$\begin{aligned} \int d^3 k k_x k_y J(k) &= \int d^3 k k k_x k_z J(k) \\ &= \int d^3 k k k_y k_z J(k) = 0, \end{aligned} \quad (76)$$

and the solution of (73) has the form

$$\mathbf{T}(\omega) = \frac{2\pi E_0 \mathbf{e}_\ell D_0(\omega) P^*(\kappa)}{1 - (\Delta\omega_0 - i\gamma_r/2) \left[\frac{1}{\omega - \omega_0 + i\gamma/2} + \frac{1}{\omega - \omega_0 + i\gamma/2} \right]}, \quad (77)$$

where

$$\begin{aligned} \Delta\omega_0 &= \Delta\omega'_0 + \Delta\omega''_0, \\ \Delta\omega'_0 &= \frac{4}{9\pi} \frac{e^2 p_{cv}^2}{\hbar \omega_g m_0^2 \nu^2} \int_0^\infty dk k^3 |P(k)|^2 \frac{P}{\kappa - k}, \end{aligned} \quad (78)$$

$$\Delta\omega''_0 = \frac{4}{9\pi} \frac{e^2 p_{cv}^2}{\hbar \omega_g m_0^2 \nu^2} \int_0^\infty dk k^2 |P(k)|^2 \left(3 - \frac{k}{\kappa + k} \right), \quad (79)$$

$$\gamma_r = \frac{8}{9} \frac{e^2 p_{cv}^2 \omega^2 \nu}{\hbar \omega_g m_0^2 c^3} |P(\kappa)|. \quad (80)$$

If the nonresonant contribution $(\omega + \omega_0 + i\gamma/2)^{-1}$ in (77) is omitted, we obtain

$$\mathbf{T}(\omega) = \frac{2\pi E_0 \mathbf{e}_\ell D_0(\omega) P^*(\kappa)(\omega - \omega_0 + i\gamma/2)}{\omega - (\omega_0 + \Delta\omega_0) + i(\gamma + \gamma_r)/2}. \quad (81)$$

Omitting nonresonant contributions and using (81), we obtain the results (19), (20) and (22) for electric and magnetic fields at $\rightarrow \infty$ with the help of substitution

$$\begin{aligned} E^\pm(\mathbf{r}, t) &= -\frac{2}{3} \frac{E_0 e^2 p_{cv}^2 \omega^2 \nu}{\hbar \omega_g m_0^2 c^2 r} (\mathbf{e}_\ell \mathbf{e}_s^\pm) \\ &\times \int_{-\infty}^\infty d\omega \frac{\omega e^{i\kappa r - i\omega t} D_0(\omega) |P(\kappa)|^2}{\omega - (\omega_0 + \Delta\omega_0) + i(\gamma + \gamma_r)/2}. \end{aligned} \quad (82)$$

For the Pointing vector at $r \rightarrow \infty$ we obtain

$$\begin{aligned} \mathbf{S}(\mathbf{r}, t, \mathbf{e}_s) &= \frac{2}{9\pi} E_0^2 \left(\frac{e^2}{\hbar c} \right)^2 \frac{p_{cv}^4 \nu}{\omega_g^2 m_0^4 c^2} |\mathbf{e}_\ell \mathbf{e}_s^\pm|^2 \frac{\mathbf{r}}{r^3} \\ &\times \left| \int_{-\infty}^\infty d\omega \frac{\omega e^{i\kappa r - i\omega t} D_0(\omega) |P(\kappa)|^2}{\omega - (\omega_0 + \Delta\omega_0) + i(\gamma + \gamma_r)/2} \right|^2. \end{aligned} \quad (83)$$

At last, the result for the scattering section at monochromatic irradiation is as follows

$$\begin{aligned} \frac{d\sigma_\mu}{d\omega_s} &= \frac{4}{9} \left(\frac{e^2}{\hbar c} \right)^2 \frac{p_{cv}^4 \omega_\ell^2}{\omega_g^2 m_0^4 c^2} |\mathbf{e}_\ell \mathbf{e}_s^\pm|^2 \\ &\times \frac{|P(\kappa)|^4}{[\omega_\ell - (\omega_0 + \Delta\omega_0)]^2 + (\gamma + \gamma_r)^2/4}. \end{aligned} \quad (84)$$

Thus, the precise results (82) - (84) distinguish from the results of the lowest approximation on the electron-light interaction only by the replacement of ω_0 на $\omega_0 + \Delta\omega$ и γ by $\gamma + \gamma_r$.

XVII. COMPARISON OF RESULTS FOR RADIATIVE DAMPING AND AMENDMENTS TO THE ENERGY WITH RESULTS OF THE QUANTUM PERTURBATION THEORY

Let us compare expression (80) for γ_r with the expression (43) from⁵. They coincide almost. Thus, one can conclude that γ_r from (82) - (84) coincides with $\gamma_{r\eta}$ with indices $\eta = 1$ or $\eta = 2$ from⁵.

As far as concerns energy amendments, we have no a coincidence with the results of the quantum theory. The calculation shows that the energy amendment of the perturbation theory, taking into account the exciton - photon transition, must be equal

$$\Delta\omega_\eta = \frac{e^2}{4\pi^2\hbar\omega_g^2m_0^2\nu^2} \int d\mathbf{k} \sum_\mu |\mathbf{p}_{cv\eta}\mathbf{e}_{\mu\mathbf{k}}|^2 \times \int dk k^3 |P_\eta(\mathbf{k})|^2 \frac{P}{k_\eta - k}, \quad (85)$$

where $k_\eta = \omega_\eta\nu/c$.

For the exciton $\Gamma_6 \times \Gamma_7$ we use vectors $\mathbf{p}_{cv\eta}$, defined in (60). Assuming $P_\eta(\mathbf{k}) = P(\mathbf{k}) = P(k)$, we have

$$\Delta\omega_1 = \Delta\omega_2 = \frac{4}{9\pi} \frac{e^2 p_{cv}^2}{\hbar\omega_g^2 m_0^2 \nu^2} \int dk k^3 |P_\eta(\mathbf{k})|^2 \frac{P}{k_\eta - k},$$

$$\Delta\omega_3 = \Delta\omega_4 = \Delta\omega_1/2. \quad (86)$$

Comparing (86) and (78) and assuming $\omega_\eta = \omega$, we find that the RHS of (86) contains an additional factor ω/ω_g in comparison to the RHS of (78), what is incidentally. However, the energy amendment to $\Delta\omega'_0$, determined in (79), is not in agreement with the results of the perturbation theory.

XVIII. RESULTS FOR SCATTERING SECTIONS

For the scattering section on excitons $\Gamma_6 \times \Gamma_7$ in spherically symmetrical quantum dots, we obtain with the help of (84) and (69a)

$$\frac{d\sigma^{++}}{d\omega_s} = \frac{d\sigma^{++}}{d\omega_s} = \frac{1}{9} \tilde{\Sigma}_0 (1 + \cos\theta)^2,$$

$$\frac{d\sigma^{+-}}{d\omega_s} = \frac{d\sigma^{-+}}{d\omega_s} = \frac{1}{9} \tilde{\Sigma}_0 (1 - \cos\theta)^2,$$

$$\tilde{\Sigma}_0 = \left(\frac{e^2}{\hbar c}\right)^2 \left(\frac{\omega_\ell}{\omega_g}\right)^2 \frac{p_{cv}^4}{m_0^4 c^2} \frac{|P(k_\ell)|}{(\omega_\ell - \tilde{\omega}_0)^2 + \Gamma^2/4},$$

$$\tilde{\omega}_0 = \omega_0 + \Delta\omega, \Gamma = \gamma + \gamma_r, k_\ell = \frac{\omega_\ell \nu}{c}, \quad (87)$$

the superscript ++ designates the polarization of incident (scattered) light $\mathbf{e}_\ell^+(\mathbf{e}_s^+)$ and so on.

Summing on polarizations of scattered light, we obtain

$$\frac{d\sigma^+}{d\omega_s} = \frac{d\sigma^-}{d\omega_s} = \frac{2}{9} \tilde{\Sigma}_0 (1 + \cos^2\theta), \quad (88)$$

where superscript $+(-)$ designates the polarization of incident light $\mathbf{e}_\ell^+(\mathbf{e}_\ell^+)$.

The total scattering section is equal pавho

$$\sigma^+ = \sigma^- = \frac{32\pi}{27} \tilde{\Sigma}_0. \quad (89)$$

Using (80) we obtain for the radiative damping that in the resonance at $\omega_\ell = \tilde{\omega}_0$ and under condition $\gamma \ll \gamma_r$

$$\sigma_{res}^+ = \sigma_{res}^- = 6\pi \left(\frac{\lambda_\ell}{2\pi}\right)^2, \quad (90)$$

where λ_ℓ is the light wave length. The result (90) is correct for any sizes of a quantum dot.

For example, using the "envelope" wave function

$$F(\mathbf{r}) = \frac{1}{2\pi R} \frac{\sin^2(\pi r/R)}{r^2} \theta(R-r), \quad (91)$$

corresponding to the low exciton energy level in a spherical quantum dot with infinitely rectangular barriers, we have

$$P(k) = \frac{2}{kR} \int_0^\pi d\kappa \sin \frac{kR_x}{\pi} \frac{\sin^2 \kappa}{\kappa}, P(0) = 1. \quad (92)$$

XIX. CONCLUSION

The problem of elastic light scattering by semiconductor quantum dots of any form, sizes and configuration is solved with the help of the semiclassical method of retarded potentials. Results are applicable for monochromatic and pulse excitations. The light refraction coefficients are assumed identical inside and outside of a quantum dot.

The electron - light interaction is taken precisely into account, i.e., all the light reradiation and absorption processes.

As an example, the differential cross section of light scattering is calculated for monochromatic light with frequency ω_ℓ on spherical quantum dot in a semiconductor of T_d class in resonance of stimulating light with exciton $\Gamma_6 \times \Gamma_7$. It is shown that at least in case of this example the exact account of electron - light interaction results only in replacement of the factor $(\omega_\ell - \omega_0)^2 + \gamma^2/4$, obtained in lowest approximation, by a factor $(\omega_\ell - \tilde{\omega}_0)^2 + (\gamma + \gamma_r)^2/4$, where $\tilde{\omega}_0 = \Delta\omega_0 + \Delta\omega$ is the exciton energy renormalized by long-distance by exchange interaction. The value γ_r is coordinated to result received with By the help of the quantum perturbation theory⁵.

Under condition $k_l R \ll 1$ (k_l is the module of the light wave vector, R is the size of a quantum dot), polarization and angular distribution of light do not depend neither on the form of quantum dot, nor from "envelope" $F(\mathbf{r})$ exciton wave function, but only from vectors $\mathbf{p}_{cv\eta}$ of nondiagonal matrix elements of exciton

momentum of excitonic state with an index η , and magnitude of scattering cross section does not depend on the sizes of quantum dot.

The results of the present work can be used for precise calculations of light absorbance by any quantum dot, which is proportional to nonradiative damping γ (see, for example,^{8,10,12?}).

At last, obtained results for the Pointing vector on

large distances from a quantum dot are applicable for the pulse irradiations. The pulse form is determined by function $D_0(\omega)$ (see(31)). It allows, for example, to describe oscillations of scattered light caused by splitting of exciton energy levels in quantum dots (compare¹⁵, where similar oscillations are predicted in light reflection and absorption by a quantum well at pulse irradiation).

-
- ¹ L.C. Andreani, F. Tassone, F. Bassani. Solid State Commun., **77**, 9, 641 (1991).
- ² L.C. Andreani, G. Pansarini, A.V. Kavokin, M.R.Vladimiriva. Phys. Rev. **B 57**, 4670 (1998).
- ³ L.C. Andreani. Confined Electrons and Photons. Ed. by E. Burstein and C. Weisbuch, Plenum Press, N.Y., 1995.
- ⁴ E. L. Ivchenko, A. V. Kavokin. Fiz. tverd. tela, **34**), N 6, with. 1815-1822 (1992).
- ⁵ S.T.Pavlov, L. I. Korovin, I.G.Lang. Proc. XIV Int. Symposium "Nanostructures: Science and Technology-2006 p. 144; cond-mat/0605650.
- ⁶ I.G.Lang, L. I. Korovin, J. A. de la Cruz-Alcaz, S.T.Pavlov. Zh. exper. teor. fiz., 2003, **123**, N 02, pp. 305-324 (JETP, 2003, **96**, N 02, pp. 268-285.); cond-mat/0212549.
- ⁷ I. G. Lang, L. I. Korovin, S. T. Pavlov, Fiz. tverd. tela, 2004, **46**, N 09, pp. 1761-1775 (Physics of the Solid State, 2004, Vol. 46, No 09, pp.1706-1718); cond-mat/0311180.
- ⁸ L. I. Korovin, I.G.Lang, D.A.Contreras-Solorio, S.T.Pavlov. Fiz. tverd. tela, **43**), N 11, with. 2091-2099 (2001); cond-mat/0104262 [Physics of the Solid State (St. Petersburg), 2001, **43**), N 11, with. 2182-2195].
- ¹⁰ L. I. Korovin, I.G.Lang, D.A.Contreras-Solorio, S.T.Pavlov. Fiz. tverd. tela, **44**, N 9, with. 1681-1689 (2002) [Physics of the Solid State (St. Petersburg), 2002, **44**, N 9, pp. 1759-1768]; cond-mat/0203390.
- ¹⁰ L. I. Korovin, I.G.Lang, S.T.Pavlov. Fiz. tverd. tela, **44**, N 9, with. 1681-1689 (2006) [Physics of the Solid State (St. Petersburg), 2006, **44**, N 9, pp. 1759-1768]; cond-mat/2006.
- ¹¹ S.V.Goupalov. Phys. Rev., **B 68**, 125311, (2003).
- ¹² L. I. Korovin, I.G.Lang, S.T.Pavlov. Fiz. tverd. tela, **44**, N 9, with. 1681-1689 (2006) [Physics of the Solid State (St. Petersburg), 2006, **44**, N 9, pp. 1759-1768]; cond-mat/0403519.
- S.T.Pavlov, L. I. Korovin, I.G.Lang. Proc. XII Int. Symposium "Nanostructures: Science and Technology-2004 p. 284.
- ¹³ I. E. Tamm. Bases of the theory of an electricity. Nauka, Moscow, 1966.
- ¹⁴ I. M. Tsidilkovsky. Zoned structure of semiconductors. Science, M. (1978).
- ¹⁵ D.A.Contreras-Solorio, S.T.Pavlov, L. I. Korovin, I.G.Lang. Phys. Rev. B., **62**, N 23, pp. 16815-16819 (2000); cond-mat/0002229.
- ¹⁶ The Raman scattering is determined by fluctuations of current and charge densities.
- ¹⁷ A fragmentation of contributions in (2) is made so that in the zero approximation (see (31) below) we have $\mathcal{E}(\mathbf{r}, \omega) = 2\pi\mathbf{e}_\ell E_0 e^{i\kappa z D_0(\omega)}$.
- ¹⁸ We assume averaging on comparatively short time, exceeding the period $T_\ell = 2\pi/\omega_\ell$, where ω_ℓ is the light frequency of a monochromatic irradiation or carrying frequency of a pulse irradiation.
- ¹⁹ Tilde over a letter means, as in (4), the replacement of vectors $\mathbf{p}_{cv\eta}$ by $\mathbf{p}_{cv\eta}^*$ and vice versa.